



Inhibitors in children and in adults

Why they are different and what do they have common?



Jan Blatny, MD, PhD

Petr Smejkal, MD, PhD

**Depts. of Paediatric/
Clinical Haematology
Centre for Thrombosis and
Haemostasis
University Hospital Brno,
Czech Republic**



evropský
sociální
fond v ČR



EVROPSKÁ UNIE



MINISTERSTVO ŠKOLSTVÍ,
MLÁDEŽE A TĚLOVÝCHOVY



OP Vzdělávání
pro konkurenceschopnost

INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ



Introduction

Inhibitors in haemophilia



- The development of inhibitors is one of the most serious complications of the treatment in patients with haemophilia
 - Brackmann et al., Eur J Haematol Suppl, 1998
- They develop in 4-20% of HA patients with the percentage rising up to 52% in certain populations
 - Manno, Haemophilia, 1999
- Less frequent in HB: 1,5-3% only –“orphan disease”
 - Lack of useful evidence on treatment outcome predictors, risk factors, diagnosis and treatment
 - Frequent allergic reactions, nephrotic syndrome
 - Di Michele, Br J hematol, 2007

How to treat inhibitor patients ?



- **ITI is the first-line treatment in children**
 - Commenced during 1st year after inhibitor development
 - In adults ITI applied remarkably less often
 - Auerswald et al., Haemophilia 2004
- **Bleeding prophylaxis prior to, during (and perhaps also outside) ITI**
 - rFVIIa
 - aPCC
- Long-time/life-long on demand (OD) treatment with:
 - rFVIIa
 - aPCC
 - Porcine FVIII (seldom in these days)
 - High dose of FVIII
 - Auerswald et al., Haemophilia 2004
 - Brackmann et al., Blood Coagul Fibrinolysis, 2000
 - Morfini et al., Haemophilia 2007




Inhibitors in children
ITI – first line treatment

ITI – three main regimens (HA)



- Malmo
 - Cost effective
 - Relatively quick
 - Uses immune adsorption and perhaps immune suppression
 - Not the “treatment of choice” in these days
- Bonn
 - Relatively expensive
 - May take up to 3 years
 - Recommended especially for high titres inhibitors
- Low(er) dose regimen
 - Relatively cheap
 - Effective enough
 - May take long time, but seldom with complications
 - CVL less imperative
- In average
 - Effective in up to 80% of cases
 - Di Michelle et al, Thromb Haemost, 2002; Smith Pathophysiol Haemost Thromb, 2002
 - Bleedings treated with by-passing agents (nowadays rFVIIa, aPCC)

ITI outcome predictors



Inhibitor titres and FVIII dose I.

Pre-ITI Historical Titters / Dose			
Historical Titer (BU)	Pre-ITI Titer (BU)	Dose (U/kg/d)	# Successes (%)
< 50	< 10	< 50	30 / 36 (83%)
		50-99	33 / 38 (87%)
		100-199	18 / 19 (95%)
		≥ 200	23 / 24 (96%)
	10 – 20	< 50	2 / 3 (67%)
		50 – 99	11 / 14 (79%)
		≥ 200	3 / 4 (75%)
	> 20	50 – 199	3 / 9 (39%)
		≥ 200	1 / 3 (33%)

*Marianni and Kroner; Immune Tolerance Study Group (ITSG)
Haematologica; 2001;86(11):1186-93*

ITI outcome predictors



Inhibitor titres and FVIII dose II.


Pre-ITI and Historical Titters / Dose			
Historical Titer (BU)	Pre-ITI Titer (BU)	Dose (u/kg/d)	# Successes (%)
> 200	< 10	< 50 – 199	5 / 11 (45%)
		≥ 200	7 / 7 (100%)
	10 - 20	< 50 – 199	1 / 3 (33%)
		≥ 200	3 / 4 (75%)
	> 20	< 50 – 199	1 / 23 (4%)
		≥ 200	12 / 18 (67%)

*Marianni and Kroner; Immune Tolerance Study Group (ITSG)
Haematologica; 2001;86(11):1186-93*

Duration of successful tolerance is dependant of pre-ITI inhibitor level



Analysis of data from the North American
Immune Tolerance Registry (n=128).
Between-group p=0.02.



DiMichele DM et al., Thromb Haemost 2002; 87:52-57.

ICP recommendations – When to start?



The International Consensus Panel (ICP)

Donna DiMichele, Chair

Charles Hay
Thiery Lambert
David Lillicrap
Steve Pipe
Mark Reding
Chantal Rothschild

Keith Hoots
Johannes Oldenburg
Pia Petrini
Mike Recht
Georges Rivard
Elena Santagostino

DiMichele DM et al., Haemophilia. 2007;13 Suppl 1:1-22



Preferred start of ITI if titre < 10 BU

When to start?

Wait until Inh titre < 10BU

- Pre ITI treatment of inh patients with by-passing agents
- Recommended : rFVIIa (90-270µg/kg/d)

Start ITI regardless of inh titre (<10 BU) if

- Waiting period > 1-2 years
- Severe/life-threatening bleeds occurs

NB: Both (aPCC and rFVIIa) recommended if bleeding occurs

Use of rFVIIa to optimise conditions for ITI



“The exclusive use of rFVIIa in acute bleeding episodes prior to commencing ITI is an effective method of decreasing inhibitor titre, thereby optimizing conditions for ITI”

Brackmann H, et al., Bood Coagul Fibrinolysis 2000; 11(Suppl 1):S39-44

I - ITI study



- Compared two ITI strategies (in HA)
 - HD (200IU/kg daily) vs. LD (50IU/kg thrice/week)
 - In **good risk**, severe, high titer HA inhibitor pts.
- **No difference in success rate**
 - (24/58 LD vs. 22/57 HD, $p=0,909$)
 - **Shorter** time to achieve neg. titer ($p=0,027$), normal recovery ($p=0,002$) and tolerance ($p=0,116$ NS) **in HD group**
- **Peak inhibitor titers correlated inversely with success rate**
 - Historical peak ($p=0,026$), on-ITI peak ($0,002$)
 - **ONLY on-ITI peak predicted** outcome ($p=0,002$)
- LD subjects bled more often ($p=0,0019$, OR 2,2)

Inhibitors in Haemophilia B



- Rare & less understood disease, mostly in severe HB patients
- High morbidity, difficult to treat
- Severe allergic reactions
 - Difficult to treat with aPCC (contains FIX)
 - rFVIIa only available for bleeding treatment
 - Effective also for prophylaxis
 - » Warrier et al. Blood Coag Fibrinol 1998, Vox Sang 1999
 - » Hay et al. BJH 2006
 - Nephrotic syndrome might complicate the disease
- ITI in HB to be considered **carefully**

ITI in Haemophilia B



- Poor overall success rate (25%)
- High risk of complication
 - Allergy, anaphylaxis, nephrotic syndrome
- Need for immune suppression to decrease risk of complications and increase success rate
 - E.g. Mycophenolate mofetil (MMF) + dexamethasone (DEXA) + IvIG
 - » Wermes et al. Blood 2000
 - » Klarmann et al. Haemophilia 2008
- Low iFIX titre before the ITI start and some way of bleeding prophylaxis is favourable

Question 1



- What will you prefer for the treatment of 2 yrs old severe HA boy with HR inhibitors?
 - A) Increase the dose of FVIII
 - B) ITI protocol
 - C) By-pass medication on demand
 - D) Prophylaxis with By-pass, but no ITI

Question 2



- Which ITI regimen will you choose for 3yrs old HA boy with peak inhibitor titre 450 BU and pre-ITI titer 15 BU?
 - A) I will not go for ITI as this a poor risk patient
 - Will continue on By-pass Tx/Prophy
 - B) I will choose LD regimen
 - C) I will go for HD regimen

Question 3



- Do you have experience with bleeding prophylaxis with By-pass medications?
 - Either within or without ITI
- A) Yes
- B) No



Inhibitors in adults

Registries


– results and success prediction



Registry	N	Dose of FVIII IU/kg/D	success (%)	Time to remission (months)	Prediction of success (P < 0,05)					
					Age at ITI start	Inhibitor titre			dose	FVIII mutation
						pre ITI	hist. peak	on ITI		
IITR	314	<50 - ≥200	51	10.5 Median	+	+	+		+	
NAITR	164 128 HR	50 – 200	63 HR	16.3 Mean		+	+	+	+	
GITR	126 104 HR	≥ 200	76 HR	7.6-15.5 Mean			+			
Spain	38	100 - 200	63 HR	9.85 Median		+	+		+	
PROFIT	103 86 HR	median 100	53 HR	8 Median		+		+		+
										47% 81%

*Franchini M, JTH 2011: 10.1007/s11239-011-6243-3

Prognostic factors of ITI success: Age and time from inhibitor detection to ITI



- **age:**

- < 5 years

**Hay, Pathophysiol Haemost Thromb 2002: (Suppl 1) 19-21*

- < 20 years:

success:

- < 5 years:

70% (38/54)

- 5 – 10 years

73% (33/45)

- 11 – 20 years

78% (36/46)

- > 20 years

40% (19/47)

**Mariani G, Semin in Thromb Hemost 2003: 69-75*

- **Interval between dg of iFVIII and start of ITI < 5 years**

- IITR (1997) success

> 70%

**Mariani G, Haematologica 2001: 1186-93*

- + iFVIII pre-ITI < 10 BU / ml

- + dose of FVIII > 100 IU / kg / D

- 65% tolerance till 1 year

- Without these three criteria:

- 65% success rate was not achieved for at last 31. months

**Ananyeva, Blood Coagul and Fibrinolys 2004: 109-124*


ITI results in adult patients from the Spanish registry



Patient	Inhibitor titre (BU/ml)		Time from inhibitor dg. to ITI (months)	Age at start of ITI (years)	FVIII (1xFIX) IU/kg/ day	Time to inhibitor elimination (months)	Inhibitor after ITI (BU/ml)
	max	pre-ITI					
1	16	16	1	22	100	2	0
2	41	-	-	21	50	3	0
3	22	8	60	25	50	4	0
4	76	17	192	32	200	4	0
5	121	7	60	57	50	7	0
6	36	2	108	35	200	8	0
7	192	22	228	39	Malmö	4	179
median	41	12	84	32	75	4	CR 6/7 (86%)
1-38 median	67	11	25	7	140	10	CR 26/38 (68%)

**Haya S. Haemophilia 2001;7:154-159*

ITI results in adult patients with VWF/FVIII: ≥ 1 poor prog f.
(age >6 years, >1 year from inhibitor development, >200 BU
hist. peak, > 10 BU pre-ITI, previously failed ITI)



Patient	Inhib. titre (BU/ml)		Time from inhibitor dg. to ITI (years)	Age at start of ITI (years)	FVIII IU/kg/day	ITI duration (months)	Inhibitor after ITI (BU/ml)
	max	pre-ITI					
1	18	5	6	45	3x weekly 50	11	< 0,5
2	50	6	18	33	3x weekly 50	21	1,22 - PR
3	30	5	13	23	3x weekly 50	4	< 0,5
4	128	6	22	23	100	33	2,2 - PR
5	737	4	18	53	3x weekly 50	25	< 0,5
6	110	5	33	54	100	12	< 0,5
7	93	2	14	24	3x weekly 50	9	< 0,5
8	45	5	1	36	3x weekly 50	12	70 - Failure
9	54	14	10	31	100	12	< 0,5
median	54	5	14	33	21,5	12	CR 6/9 (67%)
1-17							CR 9/17 (53%)


ITI results in adult patients (association of clinical outcome with inhibitor epitope profile)



Patient	Inhibitor titre (BU/ml)		Time from inhibitor dg. to ITI (months)	Age at start of ITI (years)	FVIII IU/kg/ week	ITI duration (months)	Inhibitor after ITI (BU/ml)
	max.	pre-ITI					
1	150	8	210	20	300	9	< 0,6
2	89	4	4	37	150	28,5	< 0,6
3	288	1,4	207,5	22	300	16,2	< 0,6
median	150	4	207,5	22	300	16,2	CR 3/3 (100%)
1-7 median	89	5	16	8,2	700	22,75	CR 5/7 (71%) PR 2/7 (29%)

*Gringeri A. Haemophilia 2008;14:295-302

Cost of care analysis for ITI in adult patients: comparison with rFVIIa ObsITI study: Russian experience



	No.of bleeds per year median	Dose of rFVIIa for one bleeding median	ITI dose of FVIII	CR time to CR median	Cost 10 years preITI rFVIIa (mil. €)	Cost 10 years: ITI + profyl (mil. €)	Saved during 10 years (mil. €)	Break-even point ITI + FVIII prophyl. vers. 10 years on demand rFVIIa
LR n = 3	26	4 x 96 µg/kg	50-100 U/kg daily to alter d.	3 / 3 9 months	6.4	2.2	4.2	1,2 year
HR n = 7	22	4x 104 µg/kg	100-150 U/kg á 12 h	5 / 7 10 months	3.8	2.8	1	6,8 years

* Zozulya N. WFH Congress 2010

ITI results in adult patients - Hungary



No	age (years) mean	Inhibitor titre (BU/ml)		FVIII dose (IU/kg/day) mean	Time to inhibitor elimination (months)	success
		maxim. mean	pre-ITI mean			
7	31,8	139,8	22,6	150	5,4	5/7 (71%)

**Nemes L.4th Inhibitor Workshop for Opinion Leaders in Hemophilia, Dubrovnik 2006*

ITI results in adult patients with inhibitor: – rate of success



- IITR 2001:
 - **19/47 (40%)**

- Spanish registry + Gringeri 2007:
 - $6/7 + 6/9 = \mathbf{12/16 (75\%)}$

- + Gringeri 2008 + ObsITI:
 - $+3/3 + 5/7 = \mathbf{20/26 (77\%)}$
- + Nemes 2006:
 - $+ 5/7 = \mathbf{25/33 (76\%)}$

- In total: IITR + Spanish registry + Gringeri 2x + Nemes:
 - **44/80 (55%)**

Cost-effectiveness of ITI:
patient 75 kg (max. titr <200 BU, pre-ITI < 10-20 BU)



Daily dose of FVIII	FVIII dose per year	FVIII dose per year per 75 kg	Cost of FVIII per year	Cost of FVIII per 18 months of ITI
100 IU/kg	36 500 IU/kg	2 737 500 IU	27 375 000 CZK	41 062 500 CZK
150 IU/kg	54 750 IU/kg	4 106 250 IU	41 062 500 CZK	61 593 750 CZK
200 IU/kg	73 000 IU/kg	5 475 000 IU	54 750 000 CZK	82 120 000 CZK
300 IU/kg	109 500 IU/kg	8 212 500 IU	82 120 000 CZK	123 187 500 CZK

rFVIIa 300 µg/kg per bleeding episode:

Bleeds / year	rFVIIa per year	rFVIIa/year per 75 kg	Cost of rFVIIa per year	Cost of rFVIIa per 10 years	Saved after ITI per 10 years
5	1 500 µg/kg	112 500 µg	1 935 000 CZK	19 350 000 CZK	1 800 000 CZK
10	3 000 µg/kg	225 000 µg	3 870 000 CZK	38 700 000 CZK	21 150 000 CZK
20	6 000 µg/kg	450 000 µg	7 740 000 CZK	77 400 000 CZK	59 850 000 CZK

After ITI prophylaxis FVIII 45 IU/kg/week:

FVIII per year	FVIII / year/ 75 kg	Cost of FVIII per year	Cost of FVIII per 10 years
2 340 IU/kg	175 500 IU	1 755 000 CZK	17 550 000 CZK

Question 4



- What will you prefer for the treatment of 30 yrs old severe HA man with HR inhibitors persistent from his childhood?
 - A) ITI protocol
 - B) ITI protocol according to bleeds frequency
 - C) By-pass medication on demand
 - D) Prophylaxis with By-pass, but no ITI

Question 5



- Which ITI regimen will you choose for this 30yrs old HA man with peak inhibitor titer 450 BU and present inhibitor titre 15 BU?
 - A) I will not go for ITI as this a poor risk patient
 - Will continue on By-pass
 - B) I will choose LD regimen
 - C) I will go for HD regimen
 - D) I will postpone the initiation of ITI until the inhibitor titre has dropped to < 10 BU
 - After that I will go for HD ITI

Question 6



- What will you prefer for the treatment of 40 yrs old severe HA man with new diagnosed LR inhibitors?
 - A) ITI protocol
 - B) high dose of FVIII concentrate
 - C) By-pass medication on demand
 - D) ITI protocol only if:
 - bleeding cannot be treated with FVIII

Question 7



- Which ITI regimen will you choose for this 40yrs old HA man with LR inhibitor but unsuccessful replacement therapy with FVIII?
 - A) I will choose LD regimen
 - B) I will go for HD regimen
 - C) I will go for Malmö protocol
 - D) I will use immune suppression only

Thank you!

